

<b>Fig. 1.1:</b> Different state of Matter.....	3
<b>Fig. 1.2:</b> Major discoveries in the history of Liquid Crystals .....	4
<b>Fig. 1.3:</b> General Classification of Liquid Crystals.....	6
<b>Fig. 1.4:</b> Lyotropic Liquid Crystals.....	7
<b>Fig. 1.5:</b> Representation of phase transitions for enantiotropic mesophase.....	8
<b>Fig. 1.6:</b> Representation of phase transitions for Monotropic mesophase.....	9
<b>Fig. 1.7:</b> Representation of a calamitic liquid crystals; ( $l > b$ ).....	9
<b>Fig. 1.8:</b> General template for calamitic LC molecules.....	10
<b>Fig. 1.9:</b> General template for discotic LCs and Different types of discotic LCs.....	11
<b>Fig. 1.10:</b> General template for bent-core LC compounds.....	12
<b>Fig. 1.11:</b> Orientation of the molecules in Smectic A and Smectic C phase and its respective textures .....	13
<b>Fig. 1.12:</b> Orientation of the molecules in nematic phase along its director ( $n$ ) and its respective textures (a) nematic droplets (b) schlieren structures.....	14
<b>Fig. 1.13:</b> Orientation of the molecules in nematic phase along its director ( $n$ ) and its respective textures (a) Chiral nematic phase (b) schlieren structures.....	15
<b>Fig. 1.14:</b> Various instruments for characterizing Mesomorphic properties: (a) Polarized Optical Microscopy (POM), (b) Differential Scanning Calorimetry (DSC), and (c) X-ray Diffraction (XRD).....	17
<b>Fig. 1.15:</b> Different Applications of Liquid Crystals.....	18
<b>Fig. 1.16:</b> Odd-even effect in the liquid crystal dimers.....	20
<b>Fig. 1.17:</b> Schematic representation of various types of dimers (a) Symmetric linear calamitic LC dimers; (b) Non-symmetric linear calamitic LC dimers; (c) Symmetric H-shaped calamitic LC dimers; (d) Non-symmetric H-shaped calamitic LC dimers; (e) T-shaped, (f) U-shaped, (g) O-shaped symmetric and non-symmetric LC dimers; (h) Symmetric discotic LC dimers; (i) Non-symmetric discotic LC dimers; (j) Calamitic-Discotic LC dimer (k) end to end connected bent core dimers (l) lateral-lateral	

connected bent core dimers (m, n) apex apex connected bent core dimers (o) side by side connected bent core dimers (p) Non-symmetric calamitic-bent dimer (q) Non symmetric bent core dimers (r) Non symmetric discotic-bent dimer.....	21
<b>Fig. 2A.1:</b> Some examples of Symmetrical LC dimers containing azo linkages.....	29
<b>Fig. 2A.2:</b> Reports of Symmetrical LC dimers containing azomethine linkage.....	30
<b>Fig. 2A.3:</b> DSC thermogram of (a) <i>I-4</i> and (b) <i>II-4</i> during heating and cooling cycles.....	37
<b>Fig. 2A.4:</b> Optical microphotograph of <i>I-4</i> (a) Crystalline solid (b) Cr-N transition at 218 °C on heating (c) Schlieren texture of nematic phase at 220 °C (d) N-Iso transition (e) Isotropic liquid at 228 °C (f) On cooling, Nematic droplets at 226 °C (g) coalescence of nematic droplets (h) Schlieren Texture at 224 °C (i) N-Cr transition at 199 °C.....	38
<b>Fig. 2A.5:</b> Optical microphotograph of <i>II-4</i> (a) Crystalline solid (b) Cr-N transition at 205 °C on heating (c) Schlieren texture of nematic phase at 207 °C (d) N-Iso transition 234 °C (e) Isotropic liquid at 236 °C (f) On cooling, Nematic droplets at 233 °C (g) Schlieren Texture at 232 °C (h) N-Cr transition at 202 °C (i) Crystalline state at 200 °C.....	39
<b>Fig. 2A.6:</b> Line graph of (a) <i>I-n</i> and (b) <i>II-n</i> showing relationship between flexible spacer and transition temperatures. Bar graph of (c) <i>I-n</i> and (d) <i>II-n</i> showing nematic mesophase stability of the prepared dimers in the cooling cycle.....	40
<b>Fig. 2A.7:</b> Energy-minimized Space-filling model (MM2 models derived from ChemBio 3D Ultra 14.0 software) structures of dimers showing the effect of flexible spacer.....	41
<b>Fig. 2A.8:</b> TG curve for prepared dimers (a) <i>I-6,12</i> and (b) <i>II-6,12</i> .....	44
<b>Fig. 2A.9:</b> Optimized molecular structures of all the prepared dimers of both Series <i>I-n</i> and <i>II-n</i> .....	44
<b>Fig. 2A.10:</b> Theoretical and Experimental infrared spectrum of dimers <i>I-4</i> and <i>II-4</i> predicted by using B3LYP/6-31G level of theory.....	45
<b>Fig. 2A.11:</b> ESP contours of the dimers <i>I-n</i> and <i>II-n</i> predicted by using B3LYP/6-31G level of theory.....	47

<b>Fig. 2A.12:</b> FMOs of the dimers <i>I</i> -n and <i>II</i> -n calculated by using B3LYP/6-31G level of theory.....	49
<b>Fig. 2A.13:</b> Absorption spectra for <i>I</i> -4 (a) with various UV exposure times and a 365 nm filter (b) Plot as a function of time during UV illumination ( <i>E-Z</i> ) and thermal back relaxation from photo stationary state ( <i>Z-E</i> ).....	51
<b>Fig. 2B.1:</b> Mesogenic compounds possessing naphthalene moiety.....	103
<b>Fig. 2B.2:</b> DSC Plot of compound NpAzO4Br.....	106
<b>Fig. 2B.3:</b> Optical photomicrographs of NpAzO4Br (a) Crystalline state (b) Cr-Iso at 113 °C (c) nematic droplets of Iso-N transition at 110 °C (d) schlieren texture at 108 °C.....	107
<b>Fig. 2B.4:</b> Bar graph of transition temperatures against no. of carbon of NpAzOnBr showing thermal stability of the mesophase in heating and cooling cycle.....	108
<b>Fig. 2B.5:</b> (a) SE data of Wavelength vs. Refractive index ( <i>n</i> ) and extinction coefficient ( <i>k</i> ), and (b) Wavelength vs. Real ( $\epsilon_1$ ) and imaginary ( $\epsilon_2$ ) part of dielectric constant for the NpAzO6Br.....	111
<b>Fig. 2B.6:</b> TGA plot for the compounds NpAzO4Br and NpAzO6Br.....	112
<b>Fig. 2B.7:</b> Optimized Molecular Structures of Compounds NpAzOnBr.....	112
<b>Fig. 2B.8:</b> FMO Contours of Compounds NpAzOnBr.....	114
<b>Fig. 2B.9:</b> ESP contours of Compounds NpAzOnBr.....	117
<b>Fig. 2B.10:</b> (a) Theoretical molecular structure of NpAzO6Br, atoms labelling and identification of their rings (b) Experimental infrared spectrum of NpAzO6Br compared with the corresponding predicted for the free base by using B3LYP/6-31G level of theory .....	118
<b>Fig. 2B.11:</b> Absorption spectra for NpAzO6Br (a) with various UV exposure times and a 365 nm filter (b)Thermal back relaxation process.....	120
<b>Fig. 2B.12:</b> (a) First-order plots for <i>E-Z</i> and (b) <i>Z-E</i> photoisomerization.....	121

<b>Fig. 2B.13:</b> Plot of NpAzO6Br as a function of time during UV illumination ( <i>E-Z</i> ) and during thermal back relaxation from photo stationary state ( <i>Z-E</i> ).....	121
<b>Fig. 2B.14:</b> Experimental energy gap obtained by Tauc plot for NpAzO6Br.....	122
<b>Fig. 3.1:</b> Some examples of Unsymmetrical LC dimers containing azo linkages.....	150
<b>Fig. 3.2:</b> Some examples of Unsymmetrical LC dimers containing azomethine linkages.....	151
<b>Fig. 3.3:</b> DSC thermogram of Dimers (a) <i>III-6</i> and (b) <i>IV-6</i> .....	157
<b>Fig. 3.4:</b> Optical microphotographs of dimer <i>III-6</i> (a) Cr-N transition at 172 °C during heating, (b) Schlieren texture at 174 °C, (c) N-Iso transition at 239 °C, (d) Nematic droplets at 235 °C upon cooling, (e) Schlieren texture at 233 °C, (f) N-Cr transition at 160 °C.....	158
<b>Fig. 3.5:</b> Optical microphotographs of dimer <i>IV-6</i> (a) Crystalline state at 156 °C during heating, (b) Schlieren texture at 160 °C, (c) N-Iso transition at 237 °C, (d) Nematic droplets at 234 °C upon cooling, (e) Schlieren texture at 233 °C, (f) N-Cr transition at 125 °C.....	159
<b>Fig. 3.6:</b> (a and b) Bar graphs depicting mesophase stability (c and d) Line graphs illustrating transition temperatures in heating and cooling cycle for <i>III-n</i> .....	160
<b>Fig. 3.7:</b> (a and b) Bar graphs depicting mesophase stability (c and d) Line graphs illustrating transition temperatures in heating and cooling cycle for <i>IV-n</i> .....	161
<b>Fig. 3.8:</b> Bar graph showing mesophase stability of dimers in heating cycle for <i>Series III-n</i> and <i>IV-n</i> .....	162
<b>Fig. 3.9:</b> Geometrical difference between the structure for (a) <i>III-6</i> and (b) <i>IV-6</i> shown by its planar structure .....	162
<b>Fig. 3.10:</b> TGA plot for <i>III-6</i> and <i>IV-6</i> .....	164
<b>Fig. 3.11:</b> Optimized molecular structures of dimers of <i>Series III-n</i> and <i>IV-n</i> .....	165
<b>Fig. 3.12:</b> FMOs for the dimers of <i>Series III-n</i> and <i>IV-n</i> (Values of HOMO, LUMO and $\Delta E$ are in a.u.) .....	167

<b>Fig. 3.13:</b> Space-filling model of (a) <i>III-3</i> (Odd membered dimer) and (b) <i>III-4</i> (even membered), showcasing the spacer in the all-trans conformation.....	169
<b>Fig. 3.14:</b> MEP contours of the dimers of <i>Series III-n</i> and <i>IV-n</i> .....	170
<b>Fig. 3.15:</b> Comparison of vibrational frequency obtained from experiment and calculated at B3LYP-6-31G level theory (gas phase) for <i>III-6</i> and <i>IV-6</i> .....	171
Fig. 3.16: Photoisomerization in <i>III-6</i> .....	172
<b>Fig. 3.17:</b> Absorption spectra for <i>III-6</i> (a) with various UV exposure times. The UV illumination of 0 sec ~ No UV (b) Thermal back relaxation process .....	173
<b>Fig. 3.18:</b> Tauc plot for <i>III-6</i> showing a comparison between the experimental and calculated energy band gap for both (a) <i>trans</i> and (b) <i>cis</i> form .....	174
<b>Fig. 3.19:</b> (a) $\pi$ - $A_m$ isotherm of <i>trans</i> and <i>cis-III-6</i> monolayer at 298 K and (b) Variation in the compressibility moduli ( $ E $ ) with mean molecular area ( $A_m$ ) for the Langmuir monolayer of <i>trans</i> and <i>cis- III-6</i> at 298 K.....	177
<b>Fig. 3.20:</b> BAM images of the <i>trans-III-6</i> monolayer at (a) Liquid Expanded (LE) phase, (b) Liquid expanded-liquid condensed co-existence region, (c) liquid condensed phase and (d) collapsed state at 298 K.....	177
<b>Fig. 3.21:</b> Variation in the surface pressure as the function of time after monolayer of <i>trans-III-6</i> shined with UV light for different temperatures.....	178
<b>Fig. 3.22:</b> Variation in the surface pressure as the function of time after monolayer of <i>trans-III-6</i> shined with UV light for different temperatures. The red line corresponds to the fit equation 1.....	179
<b>Fig. 3.23:</b> The variation in rate constants for photo and thermal isomerisation for Langmuir monolayer of <i>III-6</i> molecule with respect to (a) temperature (b) as a function of $\frac{1}{T}$ .....	180
<b>Fig. 4.1:</b> Some examples of Unsymmetrical Cyanobiphenyl dimers .....	233
<b>Fig. 4.2:</b> Unsymmetrical dimers with Cyanobiphenyl and azo moiety.....	234
<b>Fig. 4.3:</b> Reports on Unsymmetrical dimers with Cyanobiphenyl and azomethine/ Schiff base moiety.....	234

<b>Fig. 4.4:</b> DSC thermogram of Dimers (a) <i>V-6</i> and (b) <i>VI-6</i> .....	241
<b>Fig. 4.5:</b> Optical microphotographs of dimer <i>V-6</i> (a) Nematic phase at 178 °C during heating, (b) Schlieren texture at 181 °C, (c) N-Iso transition at 225 °C, (d) Nematic droplets at 222 °C upon cooling, (e) Schlieren texture at 218 °C, (f) N-SmC transition at 164 °C, (g) broken focal conic texture of SmC phase at 162 °C, (h) SmC-Cr transition at 148 °C, (i) Crystalline state at 143 °C.....	242
<b>Fig. 4.6:</b> Optical microphotographs of dimer <i>VI-6</i> (a) Nematic phase at 174 °C during heating, (b) Schlieren texture at 179 °C, (c) N-Iso transition at 223 °C, (d) Nematic droplets at 220 °C upon cooling, (e) Coalescence of nematic droplets at 217 °C, (f) Schlieren texture transition at 161 °C, (g) N-SmC at 158 °C, (h) focal conic texture of SmC phase at 154 °C, (i) SmC-Cr transition at 142 °C.....	243
<b>Fig. 4.7:</b> Line graph of the transition temperature vs. no. of carbons in flexible spacer for dimers of (a) <i>Series V-n</i> and (b) <i>VI-n</i> . Bar graph showing mesophase stability for dimers of (c) <i>Series V-n</i> and (d) <i>VI-n</i> .....	245
<b>Fig. 4.8:</b> Bar graph showing mesophase length of dimers in heating cycle for <i>Series V-n</i> and <i>VI-n</i> .....	246
<b>Fig. 4.9:</b> Geometrical difference between the structure for (a) <i>V-6</i> and (b) <i>VI-6</i> shown by its planar structure .....	247
<b>Fig. 4.10:</b> TGA graph for <i>V-6</i> and <i>VI-6</i> .....	248
<b>Fig. 4.11:</b> Optimized molecular structures of dimers of <i>Series V-n</i> and <i>VI-n</i> .....	249
<b>Fig. 4.12:</b> FMOs for the dimers of <i>Series V-n</i> and <i>VI-n</i> (Values of HOMO, LUMO and $\Delta E$ are in a.u.).....	251
<b>Fig. 4.13:</b> MEP contours of the dimers of <i>Series V-n</i> and <i>VI-n</i> .....	253
<b>Fig. 4.14:</b> Space-filling model (a) <i>V-3</i> (Odd membered dimer) and (b) <i>V-4</i> (even membered) (c) <i>VI-3</i> (d) <i>VI-4</i> showing spacer in all-trans conformation.....	254
<b>Fig. 4.15:</b> Comparison of vibrational frequency obtained from experiment and calculated at B3LYP-6-31G level theory (gas phase) for <i>V-6</i> and <i>VI-6</i> .....	255
<b>Fig. 4.16:</b> Absorption spectra with respect to UV exposure time for: (a) <i>V-2</i> (b) <i>V-3</i> (c) <i>V-4</i> (d) <i>V-5</i> (e) <i>V-6</i> (f) <i>V-8</i> (g) <i>V-9</i> (h) <i>V-10</i> (i) <i>V-11</i> (j) <i>V-12</i> (k) Peak absorbance picked	

up from Fig. (a) to (j) v/s UV irradiation time. Intensity of light is 1 mW/cm <sup>2</sup> and heat filter is inserted between solution and the light to avoid any heat effect arising from the UV light.....	257
<b>Fig. 4.17:</b> Thermal Back relaxation studies after UV irradiation of <i>V-n</i> series from 2-12 depicted in “a to j”. The plot of peak absorbance Vs. time of thermal back relaxation obtained from a-j is depicted in k. One can see that within 500 minutes most of the <i>cis</i> molecules are converted back to <i>trans</i> molecules.....	258
<b>Fig. 4.18:</b> First order and second order plots of the azobenzene systems studied for <i>cis-trans</i> thermal isomerization.....	259
<b>Fig. 4.19:</b> Cartoon diagram showing the distribution of spacer length along with the chemical structure.....	260
<b>Fig. 4.20:</b> Picture depicting the situation where spacer length is fixed at <i>n</i> =2.....	260
<b>Fig. 4.21:</b> Picture depicting the situation where spacer length is fixed at <i>n</i> =6.....	261
<b>Fig. 4.22:</b> Picture depicting the situation where spacer length is fixed at <i>n</i> =12.....	262
<b>Fig. 4.23:</b> Optical storage device fabricated using unsymmetrical dimers with <i>n</i> =12, where dimers were used as the guest molecule and the liquid crystal E7 is used as the host molecule. One can see from the device that exposed region is corresponding to the UV illumination (dark state) and the masked region corresponds to that UV-protected area through a mask (dark state).....	263
<b>Fig. 4.24:</b> (a) Variation of birefringence( $\Delta n$ ) as a function of temperature ( $T-T_{NI}$ ) of <i>VI-9</i> (b) Variation of dielectric permittivity at 0.5V and 5V as a function of temperature ( $T-T_{NI}$ ) of <i>VI-9</i> at applied frequency 1KHz.....	265
<b>Fig. 4.25:</b> (a) Variation of conductivity as a function of the frequency of <i>VI-9</i> (b) Variation of the real and imaginary part of dielectric permittivity ( $\epsilon'$ and $\epsilon''$ ) as a function of the frequency of <i>VI-9</i> .....	266
<b>Fig. 4.26:</b> Variation of dielectric loss ( $\tan \delta$ ) as a function of frequency of <i>VI-9</i> at $T_{NI} = -11$ °C.....	266
<b>Fig. 5.1:</b> Examples of Unsymmetrical dimers with Cholesterol and azo moiety.....	331
<b>Fig. 5.2:</b> Unsymmetrical dimers with Cholesterol and azomethine moiety.....	332

<b>Fig. 5.3:</b> DSC thermogram of Dimers (a) <i>VII-7</i> and (b) <i>VIII-7</i> .....	339
<b>Fig. 5.4:</b> Optical microphotographs of dimer <i>VII-7</i> (a) Crystalline state at 141 °C, (b) N* phase (at 143 °C) during heating, (c) N*-Iso transition at 186 °C, (d) chiral nematic droplets at 184 °C upon cooling, (e) coalescence of droplets, (f) Pseudofocal conic texture for N* at 183 °C (g) oily streak of sheared N* phase at 181 °C (h) SmC* at 117 °C (i) SmC*-Cr transition at 89 °C.....	340
<b>Fig. 5.5:</b> Optical microphotographs of dimer <i>VIII-7</i> (a) Crystalline state at 130 °C, (b) N* phase (at 133 °C) during heating, (c) N*-Iso transition at 187 °C, (d) chiral nematic droplets upon cooling at 186 °C, (e) N* phase at 185 °C, (f) N* phase on shear at 181 °C (g) N* phase at 104 °C (h) SmC* at 102 °C (i) SmC*-Cr transition at 91 °C.....	342
<b>Fig. 5.6:</b> Optical microphotographs of the textures for the LC phases of dimers: (a) Pseudo focal-conic texture of the N* phase (at 133 °C) for <i>VIII-4</i> (b) pseudo focal-conic texture of the N* phase of <i>VIII-5</i> (T = 240°C) (c) planar texture with oily streaks- N* phase for <i>VIII-5</i> (d) N* phase of <i>VIII-9</i> (T = 240°C) (e) sheared N* phase for <i>VIII-9</i> (f) N* for <i>VII-4</i> (T = 240°C) (g) N* phase (at 133 °C) for <i>VII-9</i> (h) sheared N* phase for <i>VII-9</i> (i) SmC* at 102 °C for <i>VII-9</i> .....	343
<b>Fig. 5.7:</b> X-ray diffraction spectra of <i>VII-9</i> at different temperatures showing the transition from N* to SmC* phase.....	344
<b>Fig. 5.8:</b> Line graph of the transition temperature vs. no. of carbons in flexible spacer for dimers of <i>Series</i> (a) <i>VII-n</i> and (b) <i>VIII-n</i> . Bar graph showing mesophase stability for dimers of <i>Series</i> (c) <i>VII-n</i> and (d) <i>VI-n</i> .....	345
<b>Fig. 5.9:</b> TGA graph for <i>VII-7</i> and <i>VIII-7</i> .....	346
<b>Fig. 5.10:</b> Absorption spectra of <i>VII-7</i> (a) at different UV exposure times (b) Thermal back relaxation.....	347
<b>Fig. 5.11:</b> Plot of <i>VII-7</i> as a function of time: (a) during UV illumination showing the trans-cis isomerization process, and (b) back relaxation .....	348
<b>Fig. 5.12:</b> Absorption spectra for series <i>VII-n</i> under different conditions: without UV radiation, after exposure to short UV (254 nm), and after exposure to long UV (365 nm).....	348
<b>Fig. 5.13:</b> Optimized molecular structures of dimers <i>Series VII-n</i> and <i>VIII-n</i> .....	349

<b>Fig. 5.14:</b> FMOs for the dimers of <i>Series VII-n</i> and <i>V-III-n</i> (Values of HOMO, LUMO and $\Delta E$ are in a.u.).....	351
<b>Fig. 5.15:</b> MEP contours of the dimers of <i>Series VII-n</i> and <i>VIII-n</i> .....	352
<b>Fig. 5.16:</b> Comparison of the vibrational frequency obtained from the experiment and calculated at B3LYP-6-31G level theory (gas phase) for dimers <i>VII-7</i> and <i>VIII-7</i> .....	352