

LIST OF FIGURES

Figure No.		Page No.
Chapter 2: Molecular Magnetic Materials		
2.1	FT-IR of compounds 1/2 , 3/4 and 5	17
2.2	TG-DTA of compounds (a) 1 , (b) 2 , (c) 3 , (d) 4 and (e) 5	19
2.3	Degradation pattern of compounds (a) 1 , (b) 2 , (c) 3 , (d) 4 and (e) 5	20
2.4	Solid state absorption spectra of compounds (a) 1 , (b) 2 , (c) 3 , (d) 4 and (e) 5	22
2.5	(a) Solid CD spectra of 1 and 2 in KBr pellet, (b) Solid CD spectra of 3 and 4 in KBr pellet, (c) Solid CD spectra of 5 in KBr pellet and (d) CD spectra of Ligands (<i>R</i>)-(+)-4-methyl- α -methyl benzylamine (L1), (<i>S</i>)-(-)-4-methyl- α -methyl benzylamine (L2), (<i>R</i>)-(+)-4-chloro- α -methyl benzylamine (L3), (<i>S</i>)-(-)-4-chloro- α -methyl benzylamine (L4), (<i>R</i>)-(+)-4-bromo- α -methyl benzylamine (L5)	23
2.6	Molecular view of compound 1 (a) and compound 2 (b) having thermal ellipsoid are shown at 50 % probability	24
2.7	Molecular view of compound 3 (a) and compound 4 (b) having thermal ellipsoid are shown at 50 % probability	25
2.8	Molecular view of compound 5 having thermal ellipsoid are shown at 50 % probability	26
2.9	Left handed and right handed helix motif	29
2.10	Supramolecular Cl \cdots N interactions present along the chains to	30

Designing and Studies on Molecular Materials

	stabilize Inter helical chain-3.239 Å	
2.11	Supramolecular C-H \cdots π interactions in 3 present along the adjacent chains to stabilize Inter helical chain	31
2.12	Inter helical chain distances	31
2.13	Helical chains from supramolecular interactions	32
2.14	Plots of χ_M versus T and $\chi_M T$ versus T for 1 (a), 3 (b) and 5 (c)	35

Chapter 3: Molecular Ferroelectric Materials

3.1	Schematic illustrations for the displacive-type (I) and order disorder (II)-type ferroelectrics	43
3.2	Hair like crystal growth of AnHNO ₃ (1a) from gel	51
3.3	FT-IR of compound 1a , 1b , 2a , 2b , 3a , 3b , 4b and 5b	53
3.4	Raman spectra of compound 1a , 2a , and 3a	59
3.5	¹ H NMR spectra of (a) - 1a ; (b) - 1b ; (c) - 2a ; (d) - 2b ; (e) - 3a ; (f) - 3b ; (g) - 4a ; (h) - 5a	64-67
3.6	TG-DTA of compound 1a , 1b , 2a , 2b , 3a and 3b	68
3.7	DSC chart of compounds 1a , 1b , 2a , 2b , 3a , 3b and 5b at different temperatures	70-71
3.8	(a) PE Loop measurement of AnHNO ₃ hairs; (b) PE Loop measurement of 4-EtAnHNO ₃ hairs; (c) PE Loop measurement of 4-EtAnHNO ₃ hairs with temp.; (d) PE Loop measurement of 4-EtAnHNO ₃ hairs with re-temp	74

3.9	Solid CD spectra of compound 3P and 3M in nujol	76
3.10	Molecular view of compounds 1b (a), 2b (b), 3b-M (c), 3b-P (d), 4b (e) and 5 (f) having thermal ellipsoids are drawn at the 50% probability level	80-82
3.11	Molecular view of compounds 1a (a) and 3a (b) having thermal ellipsoids are drawn at the 50% probability level	83
3.12	Images of the KNO ₃ crystal structure (a) variable K-N-K distances parallel along the <i>c</i> -axis (b) structure consists of alternating layers of K ⁺ and NO ₃ ⁻ parallel along the <i>c</i> -axis and (c) viewed parallel to the <i>c</i> -axis	84
3.13	Structure of AnHNO ₃ along <i>c</i> -axis, showing equidistant anilinium cation and nitrate anion, similar to KNO ₃ -phase-II	85
3.14	(a) Structure of AnHNO ₃ , showing interwoven double helical chains due to hydrogen bonding between anilinium cation and the nitrate anion along <i>c</i> -axis; (b) H-bonding pattern of AnHNO ₃	86
3.15	Structure of 4-MeAnHNO ₃ along <i>c</i> -axis, showing non equidistant anilinium cation and nitrate anion in hexagonal arrangement of NO ₃ ⁻ anions, similar to KNO ₃ -phase-II: (b) Hydrogen bonding assisted anti-parallel helical chains	87
3.16	(a) Hexagonal arrangement of NO ₃ ⁻ anions showing <i>p</i> -eAnH ⁺ above/below the plane for 3a/3b ; Hydrogen bonding assisted (b) left handed (<i>M</i>) helical chains : (c)) right handed (<i>P</i>) helical chains	87

3.17	Packing of the crystal structure of along the <i>c</i> -axis; carbon and hydrogen atoms are omitted for clarity	89
3.18	Pictures of hair like crystals of compound 1a and 3a	91
3.19	Packing and symmetry element 2_1 screw axes of the crystal structure of 1a and 1b along the <i>c</i> -axis	93
3.20	Hydrogen bond angles between $N^+-O^{\ominus}---N^+$ along <i>a</i> -axis in 1a and 1b	94
3.21	Arrangement of cation and anion (a) KNO_3 and (b) 3b	95
3.22	Hydrogen bonded assisted helical chains between anilinium cation and the nitrate anion along <i>c</i> -axis (a) 1b ; (b) 2b and (c) 3b	95
3.23	Packing of the crystal structure of 3a and 3b along the <i>c</i> -axis	96
3.24	Hydrogen bond angles between $N^+-O^{\ominus}---N^+$ along <i>a</i> -axis in 3a and 3b	97
3.25	(a) Molecular view of compounds 3b-HT having thermal ellipsoids are drawn at the 50% probability level; (b) Structure of 3b-HT along <i>b</i> -axis showing similar hydrogen bonded assisted helical chains as 1b; (c) Structure of 3b-HT along <i>a</i> -axis; (d) Structure of 3b-HT along <i>c</i> -axis	99
3.26	Reversible phase transition of 3b-RT ($P2_12_12_1$) to 3b-HT ($Pbca$) showing change in rearrangement of cation and nitrate anion along <i>c</i> -axis	100

Designing and Studies on Molecular Materials

3.27	Domains walls as function of temperature (30-75 °C) by polarizing microscope. The sample was placed between two polarizers that make an angle (a) 0° (re-RT sample after cooling down to the initial stage) and (b) 45°	102
3.28	(a & c) variable temperature FT-IR of 1a ; (b & d) variable temperature FT-IR of 1b and (e) variable temperature FT-IR of KNO₃	105
3.29	(a & c) variable temperature FT-IR of 3a and (b & d) variable temperature FT-IR of 3b	108

Chapter 4: Molecular Nonporous compounds for adsorption-desorption

4.1	Molecular Accommodative Chemistry with visual colour change: Non-porous Cu(II) complex, (C ₆ H ₅ CH(CH ₃)NH ₂) ₂ CuCl ₂ (A) when exposed to HCl gas (solid-gas reaction) reversibly changes to (C ₆ H ₅ CH(CH ₃)NH ₃) ₂ CuCl ₄ , double salt (B). (A) at solid-liquid interface shows Single-Crystal to Single Crystal transformation by selectively absorbing azide anion driven macromolecular assembly to form [Cu ₃ (C ₆ H ₅ CH(CH ₃)NH ₃) ₂ (N ₃) ₆] _n (C). (B) transforms reversibly to (C)	117
4.2	Compound A₁ →Compound B₁ transformation	122
4.3	FT-IR spectra of compound A₁ and B₁	123
4.4	TG/DTA curve and degradation pattern for compound A₁ (a)	125
4.5	TG/DTA curve and degradation pattern for compound B₁ (b)	125

Designing and Studies on Molecular Materials

4.6	Molecular view of \mathbf{A}_1 (thermal ellipsoid are shown at 50 % probability)	127
4.7	Molecular view of (a) \mathbf{A}_1 and (b) \mathbf{A}_1 with space filled style denotes the absence of porosity	127
4.8	Molecular view of \mathbf{B}_1 (thermal ellipsoid are shown at 50 % probability)	127
4.9	Observed and Simulated Powder X-Ray Diffraction Patterns of \mathbf{B}_1	129
4.10	FT-IR study: Monitoring of HCl gas adsorption with time during transformation of \mathbf{A}_1 to \mathbf{B}_1	131
4.11	Powder X-ray study: Monitoring (ex-situ) of HCl gas adsorption with time during transformation of \mathbf{A}_1 to \mathbf{B}_1 . (colour of the spectra are matching with the colour of the sample during this transformation)	131
4.12	Probable mechanism for transformation of (a) 1-D chain of \mathbf{A}_1 to (d) 2-D layer of \mathbf{B}_1 through (b) benzylammonium formation, and (c) charge neutrality driven layer formation at the expense of inversion center between the layers	132
4.13	Observed (KOH treatment- \mathbf{A}_1 r) and Simulated Powder X-Ray Diffraction Patterns of \mathbf{A}_1 (Note the peak due to eliminated KCl at $2\theta=28^\circ$)	133
4.14	Compound $\mathbf{A}_R / \mathbf{A}_S \rightarrow$ Compound $\mathbf{B}_R / \mathbf{B}_S$ transformation	137
4.15	FT-IR spectra of compound $\mathbf{A}_R / \mathbf{A}_S$ and $\mathbf{B}_R / \mathbf{B}_S$	137

4.16	TG/DTA curve and degradation pattern for compounds \mathbf{A}_R (a) and \mathbf{A}_S (b)	140
4.17	TG/DTA curve and degradation pattern for compounds \mathbf{B}_R (a) and \mathbf{B}_S (b)	141
4.18	Molecular view of (a) \mathbf{A}_R and (b) \mathbf{A}_S (thermal ellipsoid are shown at 50 % probability); (c) \mathbf{A}_R with space filled style denotes the absence of porosity	142
4.19	Molecular view of (a) \mathbf{B}_R and (b) \mathbf{B}_S (thermal ellipsoid are shown at 50 % probability)	142
4.20	Solid state absorption spectra of conversion of $\mathbf{A}_S \rightarrow \mathbf{B}_S$	145
4.21	(a-c) ex-situ FT-IR monitoring of $\mathbf{A}_R/\mathbf{A}_S \rightarrow \mathbf{B}_R/\mathbf{B}_S$ in KBr pellet	146
4.22	PXRD pattern of \mathbf{B}_R obtained after keeping \mathbf{A}_R during 150 minutes in the HCl gas environment after the solid-gas interface synthesis. Wavelength: 1.5405 Å	147
4.23	PXRD time-dependent transformation of \mathbf{A}_R to \mathbf{B}_R in HCl gas environment measured at 298 K. Wavelength: 1.5405Å	147
4.24	XRPD of conversion of $\mathbf{B}_S \rightarrow \mathbf{A}_S$ (r) after KOH. (Note the peak due to eliminated KCl at $2\theta=28^\circ$)	148
4.25	Solid state CD spectra of \mathbf{A}_R , \mathbf{A}_S , \mathbf{B}_R and \mathbf{B}_S in nujol	148

4.26	Circular Dichroism study: Monitoring of HCl gas adsorption with time during transformation of \mathbf{A}_1 to \mathbf{B}_1 and desorption \mathbf{A}_S (r). (colour of the spectra are matching with the colour of the sample during this transformation)	149
4.27	Theoretical study on \mathbf{A}_S and \mathbf{B}_S	151
4.28	Different structures of ligands employed during present investigation	152
4.29	Solid state EPR spectra of \mathbf{A}_S and \mathbf{B}_S at room temperature	154
4.30	(a) DSC and (b) Thermo-chromic behavior for \mathbf{B}_1 using polarizing microscope	155
4.31	(a) DSC of \mathbf{B}_R , (b) DSC of \mathbf{B}_S and (c) Thermo-chromic behavior for $\mathbf{B}_R / \mathbf{B}_S$ using polarizing microscope	156
4.32	PE Loop measurement of \mathbf{B}_R	158
4.33	Role of molecules during self-assembly and molecular recognition, a true anchimeric assistance during crystal to crystal transformation from \mathbf{A} to \mathbf{C}	160
4.34	Crystal transformation of $\mathbf{B}_R/\mathbf{B}_S \rightarrow \mathbf{C}_R/\mathbf{C}_S$	163
4.35	Crystal transformation of $\mathbf{C}_R/\mathbf{C}_S \rightarrow \mathbf{B}_R/\mathbf{B}_S$	163

Designing and Studies on Molecular Materials

4.36	Molecular view of (a) C_R and (b) C_S ; Arrangement of the 2D brick-wall network (c) C_R and (d) C_S (H and C atoms of the amine ligands are omitted for clarity)	165
4.37	Observed and simulated XRPD pattern of C_R	166
4.38	(a) Observed and simulated XRPD pattern of C_R ; (b) Observed and simulated XRPD pattern of B_R	166
4.39	Solid state CD spectra in nujol (a) A_R and A_S ; (b) C_R and C_S	168
4.40	Solid state CD spectra in nujol (a) B_R and B_S ; (b) C_R and C_S	169

Chapter 5: Racemic Resolution in Molecular Materials

Part-I: Chloride anion assisted Resolution of chiral cobalt compounds

5.1	FT-IR spectra of compounds $1R$, $1S$, 2 and $3R$	180
5.2	TG/DTA thermogram for compounds $1R$ (a), $1S$ (b), 2 (c) and $3R$ (d)	183
5.3	Degradation pathway for compounds $1R$ (a), $1S$ (b), 2 (c) and $3R$ (d)	184
5.4	Molecular view of compounds $1R$ (a), $1S$ (b), 2 (c) and $3R$ (d) having thermal ellipsoid are drawn at the 50% probability level	188-189
5.5	2D sheet arrangement adopted by $1R/1S$ with two independent circuits A and B, propagated by N-H \cdots Cl-Co and N-H \cdots Cl hydrogen bonds	191

Designing and Studies on Molecular Materials

- 5.6** Linking of 4-F phenylethylammonium cations and tetrachloro cobaltate anions through N–H \cdots Cl hydrogen-bonded contacts to form a 1-D chain (tapes) of alternating cations and anions in compound **2** 191
- 5.7** Helical chains of compounds (a) 2-racemic P+M, (b) **1R–M**, and (c) **1S–P**. Hydrogen atoms and CoCl₄²⁻ are omitted for clarity 192
- 5.8** Adjacent helices (opposite in direction) generated by 2-fold screw symmetry operations along the b-axis in case of compounds **1R** and **3R** while opposite in case of **1S** 193

Chapter 5: Racemic Resolution in Molecular Materials

Part II: A case study of Cu-azide bridged compounds with *racemic* auxiliary ligand

- 5.9** FT-IR spectra of compounds **3/4** and **5** 202
- 5.10** FT-IR spectra of compounds **6** and **7** 203
- 5.11** (a) Compound **3(R)**, (b) Compound **5 (RS)**, (c) Compound **4 (S)**; Red circles indicate the chiral centers (d) & (e) Dihedral Angle between two ligands of compound **4 (S)**- 15.85° and **5 (RS)**- 0.00° 207
- 5.12** 1D Left handed (Λ) and Right (Δ) handed helical chains along 2₁ screw axis for **6** and **7**, respectively, showing (A) EE-azide network between Cu(II) ; (B) & (C) hydrogen bonding between N-H amine 211
- 5.13** (a) Rotation hindered due to Strong H-bonding of Ligand and azide (b) Supramolecular Cl \cdots N interactions (Cl \cdots N = 3.239 Å, blue lines) along the helical chains 211

Designing and Studies on Molecular Materials

5.14	(a) Solid CD spectra of 3 (<i>R1-R5</i>) and 4 (<i>S1-S5</i>) in nujol, (b) Solid CD spectra of compound 5 (<i>rac1- rac 10</i>) and bulk sample in nujol, (c) Solid CD spectra of 5 (<i>R1-R5</i>) and 6 (<i>S1-S5</i>) in nujol and (d) CD spectra of bulk of sample	212
------	---	-----

Chapter 6: Molecular Thermochemical Materials

6.1	FT-IR spectra of [(<i>R</i>)-(+)-1-C ₁₀ H ₇ CH(CH ₃)NH ₃]CuCl ₄ (3) and [(<i>R</i>)-(+)-2-C ₁₀ H ₇ CH(CH ₃)NH ₃]CuCl ₄ (4)	226
6.2	TG/DTA thermogram for compounds 3 (a) and 4 (b).	229
6.3	Degradation pathway for compounds 3 (a) and 4 (b)	229
6.4	DSC measurement plots for compounds 3 (a), 4 (b) at 300-450K and 3LT (c) at 200-300K	231
6.5	Thermochemical behavior for compound 3 using polarizing microscope	232
6.6	Thermochemical behavior for compound 4 using polarizing microscope	233
6.7	Representation of the ellipsoid of thermal mean square displacements of non-hydrogen atoms at 50% of the asymmetric unit of (a) compound 3 and (b) compound 4	234
6.8	View of the copper (II) coordination sphere in (a) 3LT ; (b) 3HT	237
6.9	Hydrogen bonding pattern of compound 3 at different temperature. (N-H \cdots Cl hydrogen-bonded supramolecular motif)	238

Designing and Studies on Molecular Materials

6.10	Supramolecular C-H \cdots π interactions in 3 present along the ligands	239
6.11	View of the copper (II) coordination sphere in 4HT . Thermal ellipsoids are drawn at 50% probability	240
6.12	Hydrogen bonding pattern of compound 4 at different temperatures	241
6.13	Supramolecular C-H \cdots π interactions in 4 (continuous) and 4HT (non-continuous) present along the ligands	241
6.14	Variable temperature FT-IR spectra of compound 3	245
6.15	Variable temperature FT-IR spectra of compound 4	245
6.16	Showing the solid state electronic spectra of the ligand hydrochloride (1HCl and 2HCl) and compounds 3 and 4 in nujol	247
6.17	Solid state fluorescence behavior of ligand hydrochloride (1HCl and 2HCl) and compounds 3 and 4 in nujol	248

Chapter 7: Molecular Dual Fluorescent Materials

7.1	Jablonski diagram illustrates a singlet ground electronic state (the parallel bars labeled S(0)), as well as singlet first (S(1); upper set of parallel bars, sphere representing an electron)	253
7.2	(a) Pictorial presentation of State reversal model (b) Fluorescence spectra of DMABN (1) in several solvents: (1) n-hexane, (2) dibutyl ether, (3) diethyl ether, (4) butyl chloride, and (5) acetonitrile [2]	254

Designing and Studies on Molecular Materials

7.3	The twisted intramolecular Charge transfer (TICT) model	255
7.4	(a) Fluorescence spectra of DMABN (I) in non polar solvents: (1) chloroform, (2) CCl ₄ ; (II) in polar aprotic solvents: (1) DCM, (2) ACN, (3) EtOAc, (4) Acetone; (III) in polar protic solvent: (1) Methanol. (b) Possible twisting or rotation of Donor as well as Acceptor group	259
7.5	FT-IR of compounds 1 , 2 , 3 and 4	264
7.6	¹ H NMR spectra of 4-thiocyanato aniline (1)	268
7.7	¹ H NMR spectra of N- methyl-4-thiocyanato aniline (2)	268
7.8	¹ H NMR spectra of N, N- dimethyl-4-thiocyanato aniline (3)	269
7.9	¹ H NMR spectra of 2,6-di methyl-4-thiocyanato aniline (4)	269
7.10	TG-DTA of compounds 1 (a), 2 (b), 3 (c) and 4 (d)	270
7.11	Molecular view of compounds 1 (a) and 3 (b) having thermal ellipsoid is drawn at the 50% probability level	272
7.12	Molecular view of compound 4 having thermal ellipsoid is drawn at the 50% probability level	273
7.13	Normalized absorption (Abs) and fluorescence spectra of the four thiocyanato aniline and its derivatives in chloroform (a) at 25°C	279
7.14	Normalized absorption (Abs) and fluorescence spectra of the four thiocyanato aniline and its derivatives in CCl ₄ (b) at 25°C	280

Designing and Studies on Molecular Materials

7.15	Normalized absorption (Abs) and fluorescence spectra of the four thiocyanato aniline and its derivatives in DCM (c) at 25°C	281
7.16	Normalized absorption (Abs) and fluorescence spectra of the four thiocyanato aniline and its derivatives in ethyl acetate (d) at 25°C	282
7.17	Normalized absorption (Abs) and fluorescence spectra of the four thiocyanato aniline and its derivatives in acetonitrile (e) at 25°C	283
7.18	Normalized absorption (Abs) and fluorescence spectra of the four thiocyanato aniline and its derivatives in methanol (f) at 25°C	284
7.19	FT-IR of compounds 5 and 6	288
7.20	¹ H NMR spectra of compound 5	291
7.21	¹ H NMR spectra of compound 6	291
7.22	Molecular view of compounds 5 (a) and 6 (b) having thermal ellipsoid is drawn at the 50% probability level	292
7.23	Absorbance and fluorescence spectra of the compounds 5 (a) and 6 (b) in acetonitrile at 25°C	297

Chapter 8: Interaction of Molecular Nanoparticles and CdS nanoparticles

8.1	Molecular orbital diagram of Metallo porphyrin	306
8.2	FT-IR Spectrum of compound 1	310
8.3	FT-IR Spectrum of compound Zn-1	310

Designing and Studies on Molecular Materials

8.4	1H NMR Spectra of (a) 1 and (b) Zn-1	312
8.5	Absorption spectra of (a) 1 and (b) Zn-1	314
8.6	Fluorescence study of (a) 1 and (b) Zn-1	316
8.7	Charts indicating particle size of 1NC , Zn-1NC and 3NC in solution using polyacrylamide stabilizer	321
8.8	Charts indicating particle size of 1NC (ex-situ) and 1NC (in-situ) in solution using polyacrylamide stabilizer	321
8.9	Absorption spectra of CdS and its nanoparticles in polyacrylamide	323
8.10	Normalized absorption Spectra of (a) 1 / 1-NP / 1-NC (b) Zn-1 / Zn-1NP / Zn-1NC	324
8.11	Fluorescence spectra of 3	326
8.12	Comparison between fluorescence spectra of (a) 1 and 1NP , (b) Zn-2 and Zn-1NP , (c) 1NP and 1NC and (d) Zn-2NP and Zn-1NC	327
8.13	CdS colloids surface interaction with (a) 1NC , (b) Zn-1NC	330

Annexure I: Home Built Variable Temperature based Infra Red Spectroscopy Attachment

I.1	The schematic of the FTIR attachment which has three parts namely: 1-Sample holder, 2-Heater and 3-PID based temperature controller	335
I.2	Actual setup heater assemble fitted inside the FTIR sample holder	336

Designing and Studies on Molecular Materials

I.3	Whole set up for Temperature controlled FTIR spectra sample holder	336
------------	--	-----

Annexure III: Synthesis of strong electron acceptor compounds

III.1	FT-IR spectra of compound 1 (a) and compound 2 (b)	352
III.2	¹³ C NMR spectra of compound 1 (a) and compound 2 (b)	355
