

List of Figures

Figure No.	Figure Caption	Page No.
1.1	Tris chelate complexes	02
1.2	Examples of NILs	05
1.3	Classification of NILs	06
1.4	1-(α -amino benzyl)-2-naphthol (Betti base)	11
1.5	Applications of Betti base	12
2.1	Crystal structure of <i>S</i> -Betti base hydrobromide	21
2.2	Image of Potentiostat with Electrochemical cell	29
2.3	Structure of Betti base and its derivatives	34
2.4	Circular Dichroism spectra (0.5mM) in 10mL of chloroform. (a) CD spectra of L₂ and L₃ (b) CD spectra of L₅ and L₆	36
2.5	Cyclic voltammograms of Betti bases (1mM in DCM) using 0.1 M TBAPF ₆ as the supporting electrolyte.	37
2.6	Cyclic voltammograms of L₁ (1mM in DCM) using 0.1 M TBAPF ₆ as the supporting electrolyte. (A) 0.5 V to -0.9 V (B) 1.2 V to -0.9 V and (C) 1.2 V to -0.9 V starting from 0.0 V	38
2.7	Cyclic voltammogram of the Betti base at increasing scan rate from 100 to 500 mV/sec	39
2.8	Plots of square root of scan rate vs. current (a),(b) cathodic current ; (c),(d) anodic current	40
2.9	Cyclic voltammograms of a 1 mM solution of BB in DCM at 300 mV/sec containing 0.1 M TBAPF ₆ as the supporting electrolyte.	41
2.10	CVs of L₄	43
2.11	Plots of current vs. scan rate	43
2.12	CVs of L₄ in different pH	44
2.13	CV of L₄ and L₇	44
2.14	CVs of L₇ in different pH	45
2.15	Structures of the α -alkyl amino phenol derivatives	45

2.16	E4 Plots for (a) Oxidation process (b) Reduction process of Betti base	55
2.17	E2 Plots for (a) oxidation of Betti base (b) reduction of Betti base; E3 Plots for (c) oxidation of Betti base (d) reduction of Betti base	55
2.18	E4 Plots for (a) Oxidation process at low pH of Betti base (b) Oxidation process at high pH of Betti base	56
2.19	E2 Plots for Betti base at high pH (a) oxidation (b) reduction; E3 Plots for Betti base at high pH (c) oxidation (d) reduction	56
2.20	E4 Plots for for PBB (a) Oxidation process (b) Reduction process	58
2.21	Plots for elimination functions E2 (a) oxidation of PBB (b) reduction of PBB ; plots for the elimination function E3 (c) oxidation of PBB (d) reduction of PBB	59
2.22	Plots for diffusion current I_d for PBB at high pH (a) Oxidation process (b) Reduction process	59
2.23	Plots for elimination functions E2 at high pH (a) oxidation of PBB (b) reduction of PBB ; plots for the elimination function E3 at high pH (c) oxidation of PBB (d) reduction of PBB	60
2.24	E4 Plots for DBB (a) Oxidation process (b) Reduction process	61
2.25	E4 Plots for ABB (a) Oxidation process (b) Reduction process	62
3.1	The number of papers concerning the Cu(II) complex with (blue) N,O donor ligands, (red) amino phenols and (green) amino naphthols being published within 2005-2014 (Data taken from Thompson Reuters)	73
3.2	Mono and bis(pyridyl)alkylamine ligands	74
3.3	Schematic representation of the structure of the mixed Cu(II)–phenolate/phenol complex	75
3.4	Schematic representation of $Tp^R Cu(II)(phenolate)$ complexes, where X = a exogenous, bidentate phenolate ligand ($R_1 = R_2 =$ alkyl substituents).	75
3.5	Structure of 2-Anilino-4,6-di- <i>tert</i> butylphenol and its Cu(II) complex	76
3.6	Structure of 2-[2'-(4',6'-di- <i>tert</i> butylhydroxyphenyl)]- 4,5-diphenylimidazole and its Cu(II) complex	76
3.7	Structure of tripodal amino phenol and its Cu(II) complex	77

3.8	Structure of N-Phenyl-2-amino phenol substituted ligand and its Cu(II) complexes	77
3.9	Structures of (a) benzoxazole iminosemiquinone ligand and its Cu(II) complex (b) iminophenol amidopyridine ligand and its Cu(II) complex	79
3.10	(a) TGA of Cu1 (b) TGA of Cu4 (c) TGA of Cu7 (d) TGA of Cu8 (e) TGA of Cu9	83
3.11	(a) UV spectra of Cu3 & Cu6 (b) Visible spectra of Cu3 , Cu6 & Cu7	85
3.12	Circular Dichroism spectra (0.5mM) in 10mL of chloroform (a) CD spectra of Cu2 and Cu3 (b) CD spectra of Cu5 and Cu6	86
3.13	EPR spectra of Cu(II) complexes prepared in DCM (10 mM). Spectra were collected at Room temperature, $\nu_0 = 9.1$ GHz. (a) Cu1 (b) Cu2 (c) Cu4 and Cu5	88
3.14	ORTEP diagram of Cu1	89
3.15	Proposed structure of Cu2	91
3.16	(a) and (b) cyclic voltammogram of Cu2 with increasing scan rate 10 mV/sec to 40 mV/sec; (c) plot of the cathodic peak current for Cu2 as a function of square root of scan rate in the range 10 mV to 50 mV/s. (d) plot of the anodic peak current for Cu2 as a function of square root of scan rate in the range 10 mV to 50 mV/s.	93
3.17	(a) E4 elimination plot for the oxidation of Cu2 (b) E4 elimination plot for the reduction of Cu2 .	94
3.18	Plots for elimination functions E2 (a) oxidation of Cu2 (b) reduction of Cu2 ; plots for the elimination function E3 (c) oxidation of Cu2 (d) reduction of Cu2	96
3.19	(a) cyclic voltammogram of Cu5 with increasing scan rate 100 mV/sec to 400 mV/sec; (b) plot of the cathodic and anodic peak current for Cu5 as a function of square root of scan rate in the range 100 mV to 500 mV/s.	97
3.20	(a) E4 elimination plot for the oxidation of Cu5 (b) E4 elimination plot for the reduction of Cu5 .	97
3.21	Plots for elimination functions E2 (a) oxidation of Cu5 (b) reduction of Cu5 ; plots for the elimination function E3 (c) oxidation of Cu5 (d) reduction of Cu5	98

4.1	(A) Effect of amount of oxidant on styrene conversion (B) Effect of amount of catalyst on styrene conversion.	115
4.2	Electronic spectra of (a) Cu1 in acetonitrile (b) Cu1 + TBHP in acetonitrile (c) Cu1 +TBHP + Styrene in acetonitrile after 24 h	117
4.3	(A) FTIR spectra of Cu1 (B) Overlaid FTIR spectra of Cu1 during reaction (a) Cu1 after 2 h (b) Cu1 after 4 h	119
4.4	CV of Cu1 in ACN (5 mmol)	120
4.5	The ¹ H NMR spectra of TBHP treated Betti base in CDCl ₃	122
4.6	(A) Effect of amount of oxidant on styrene conversion (B) Effect of amount of catalyst on styrene conversion	125
4.7	Chiral GC-MS analysis for epoxidation of styrene in DMSO	128
4.8	(A) FTIR spectra of Cu1 (B) Overlay FTIR spectra of Cu1 during reaction a. Cu1 after 2 h b. Cu1 after 4 h c. Cu1 after 24 h.	130
4.9	(a) CV of Cu1 in DMSO (5 mmol) (b) CV of Cu1 in ACN (5 mmol) (c) CV of Cu1 in DMSO	131
4.10	CV of Cu1 in DMSO; scan rate 10 – 80 mVs ⁻¹ ; room temperature	132
5.1	Chiral Schiff base Cu(II) complex	144
5.2	Aratani's copper catalyst used in the commercial manufacture of <i>cilastatin</i>	144
5.3	Chiral C ₂ -symmetric bisoxazoline ligand	145
5.4	First semicorrin type ligand	145
5.5	Masamune and Evans' catalysts	145
5.6	Schiff base used by Somanathan and Hake	147
5.7	(a) Cu(II) Betti base complex (b) Cu(II) Schiff base complex	148
5.8	EPR spectra of Cu(II) complexes (10 mM in DCM). Spectra were collected at room temperature, $\nu_0 = 9.1$ GHz. (a) Cu1 (b) Cu2 (c) Cu5 & Cu6	151
5.9	Two addition mechanisms for cyclopropanation reactions	152
5.10	Probable pathway for the formation of metal-carbene adduct	153